

TPC Distortions & Calibrations

estimated magnitudes and our ability to correct



STAR TPC Review
BNL, June 4-5, 2009

Outline


- Overview of the distortions
 - Description
 - Impacts of high luminosity (and aging)
- Correction techniques (calibrations)
 - Physics requirements
- Results (past and future prospects?)
- Summary

Distortions

- EM fields: non-uniformities are a reality
 - B field: small deviations (scale?), mapped, done
 - E field: surface & volume issues, static & volatile
 - The big three: (1) shorted field cage rings, (2) primary space charge, (3) gated grid ion leakage
- Electrostatics is known physics
 - Requirements: (1) model of the distortion, (2) measures/rulers (e.g. surveys, residuals) which keep pace with volatility

Field Cage Electrical Shorts

- Potential stepped from cathode to anode
- “Stripes” express potential inside the chamber




Field Cage Electrical Shorts

- Shorts have been a problem for several years now
 - Some fixed
 - Some not understood
- Worst threat comes from volatile shorting
 - Not a current problem
- Very unlikely a high luminosity issue
 - Unknown whether its an aging issue

SpaceCharge: model of charge


- HIJET model of “event shape” for 200 GeV AuAu collisions matches radial distribution of zerobias data well for much of the runs.

Radial distribution of TPC SpaceCharge



SpaceCharge: model of charge


- HIJET model of “event shape” for 200 GeV AuAu collisions matches radial distribution of zerobias data well for much of the runs.



March 1, 2004 data


SpaceCharge effect on sDCA

J. Dunlop



- All tracks go the same direction (pos. or neg.)
- Track charge independence
- Field dependence


sDCA = signed distance of closest approach




SpaceCharge effect on sDCA

J. Dunlop

Vertex-finding de-focused,
but not biased:
vertex makes a good
reference point




sDCA = signed distance
of closest approach




GridLeak Field Effects


- Modeled sheets of charge
 - Relaxation done on custom 3D grid
(plots assume Φ symmetry, but leak is 12-fold symmetry from grid shape)
 - E-field and distortion discontinuity at grid gap
- GridLeak scales as SpaceCharge!



Simulated residuals on a track





Distortion near CM



Applied GridLeak Correction

• Not perfect,
but as good as
design spec!

Distortions scale
significantly reduced!




After

Before

Projected pointing errors


- pp500 is the worst
- Run 9 was a good test
- Intermediate ions (CuCu) perhaps worse than heavy ions (AuAu, UU?)

TPC pointing error vs. luminosity (no corrections)




Pointing resolution

- Important for using inner (silicon tracking, upgrades)
- More Discussion



Momentum resolution

- Important for physics at high p_t
- Biases can be more serious than smearing
- More Discussion



Distortion Corrections

Distortion	Approximate Scale [microns]	Correction Scale [microns]
Twist (E-B alignment)	800	50
IFC Shift	100	50
Clock (East-West rotation)	800	50
Padrow 13	400	50
B field shape	800	50
Shorted Ring	2000 ^A	100 ^B
Space Charge	up to 5000 ^C	100-200 ^D
Grid Leak	up to 2500 ^C	100-200 ^D
Unknown	100??? 300???	100??? 300???

- Overall contribution to $\delta p_t/p_t \sim 1/4 - 3/4\% * p_t$ for TPC-only tracks (primary vtx, silicon help)

A. Larger (up to 5000) without compensating resistor.

B. Known to be ~400 microns in a region of the TPC not used for physics.

C. Luminosity dependent


D. Dataset dependent

CDR design was ~1%*pt

Momentum biases

- A very small bias can lead to a large effect
- h^-/h^+ with a mere $0.005 \cdot p_t^2$ bias
- More discussion

$$\frac{\text{pow}([0]+x+[1]*x*x,[2])}{\text{pow}([0]+x-[1]*x*x,[2])}$$



Calibration first steps

- Non-volatile calibrations must be completed first (e.g. internal alignment, and w.r.t. B field)
 - Necessitates low luminosity data
 - RHIC was unable to deliver this during pp500 this year.....the future?
- Field cage currents measure shorts
- sDCA and residuals tell us about the SpaceCharge and GridLeak


First steps to corrections

- Observables (sDCA) can tell you the distortion quantity (ions in the TPC due to SpaceCharge buildup + GridLeakage)
- Easy with “ideal” tracks
 - Little or no dependencies on reconstruction itself
 - Observable maps easily to distortion quantity
 - $sDCA = C * f(Z) * (SpaceCharge + GridLeak)$
 - Generally need many events for stats
 - Could be many runs for pp collisions!

First steps to corrections

- Observables (s distortion quant SpaceCharge b
- Easy with “ideal
- Little or no depe
- Observable map


Space Charge



- $sDCA = C * f(Z) * (SpaceCharge + GridLeak)$
- Generally need many events for stats
 - Could be many runs for pp collisions!

Ionization: Scalers


- Ionization is linear with scaler measures of luminosity
- Points out problem runs
- Now using 1-second averages



STAR records scaler rates on Zero Degree Calorimeters (ZDCs) and Beam-Beam Counters (BBCs)

Volatility seen during AA


- Fluctuations seen on the 1-second time scale
- More discussion



Performance Measures: sDCA

2004 AuAu at 200 GeV, all B fields


- Can't beat low luminosity, but holding steady at high luminosity:
- Spread from 5-9kHz appears roughly uniform
- No indication we can't go higher!



Performance Measures: sDCA


2005 CuCu at 200 GeV, full field

- Can't beat low luminosity, but holding steady at high luminosity:
- E-by-E method performs worse due to statistics per unit time



Performance Measures: π^-/π^+

- TPC-measure of the ratio essentially flat all the way to $p_T=12$ GeV/c !
- Central triggers (taken at high luminosity) just about as good!




O. Barannikova

The Future: Up and Up (1)

- Higher luminosities
 - Can't even do tracking across TPC without some GridLeak correction
 - How close is our model to reality?
 - Differences will amplify with increasing luminosity.
 - How will the backgrounds change/grow/quell?
 - Not clear that the shielding has removed non-collision contributions
 - pp500 has been a valuable test...

pp500: getting pretty high

- Preliminary calibration modestly successful
- No major surprises
- Non-linearity already encountered in Run 8 dAu 200 GeV.
- More Discussion



The Future: Up and Up (2)

- Higher DAQ rates
 - Increasing gating grid rates produced no notable change in SpaceCharge-like distortions(!)
 - Higher event rate *might* benefit the E-by-E approach
- Other techniques for SpaceCharge measures
 - Fixed detectors (GMT upgrade proposal)
 - Use identified pileup hits in the data (work in progress)

Back to the table...

- Analyses requirement: don't gain another $\sqrt{2}$
- What can we afford?
 - It is NOT the increasing distortions which hurt most, it is the increasing error of our understanding!
- Room to increase the error on our luminosity-dependent corrections
 - Hard to say what will happen...

Distortion	Approximate Scale [microns]	Correction Scale [microns]
Twist	800	50
IFC Shift	100	50
Clock	800	50
Padrow 13	400	50
B field shape	800	50
Shorted Ring	2000 ^A	100 ^B
Space Charge	up to 5000 ^C	100-200 ^D
Grid Leak	up to 2500 ^C	100-200 ^D
Unknown	100??? 300???	100??? 300???

"Overall contribution to $\delta p_t/p_t \sim 1/4\text{-}3/4\%$ * p_t for TPC-only tracks (primary vtx, silicon help)"

- A. Larger (up to 5000) without compensating resistor.
- B. Known to be ~ 400 microns in a region of the TPC not used for physics.
- C. Luminosity dependent
- D. Dataset dependent

Our efforts are worthwhile!

- STAR TPC has major distortions with which we have been coping for years now (Physics produced!)
- Preliminary efforts appear successful with pp500, but we expect even higher luminosities and things could get worse for us
- We have some margin for further resolution error, but even small biases are problematic



Extras

Backgrounds

- Strong evidence for collider backgrounds: calorimeter backgrounds, “straight-through” tracking, zero-bias data, sDCA azimuthal distributions
- Will shielding solve this problem for good?
- Will we need to “map” the distortions?

